Given a set *L* of (possible) locations for placing facilities and a set *D* of customers together with cost functions  $s: D \times L \to \mathbb{R}^+$ and  $o: L \to \mathbb{R}^+$  find a set of facility locations *F* together with an assignment  $\phi: D \to F$  of customers to open facilities such that

$$\sum_{f\in F} o(f) + \sum_{c} s(c, \phi(c))$$

is minimized.

In the metric facility location problem we have

$$s(c, f) \le s(c, f') + s(c', f) + s(c', f')$$
.



◆ 個 ト ◆ 三 ト ◆ 三 ト 476/491

#### **Integer Program**

min		$\sum_{i \in F} f_i \gamma_i + \sum_{i \in F} \sum_{j \in D} c_{ij} x_{ij}$		
s.t.	$\forall j \in D$	$\sum_{i\in F} \chi_{ij}$	=	1
	$orall i \in F$ , $j \in D$	$\chi_{ij}$	$\leq$	${\mathcal{Y}}_i$
	$\forall i \in F, j \in D$	$x_{ij}$	$\in$	$\{0, 1\}$
	$\forall i \in F$	${\mathcal Y}_i$	$\in$	$\{0, 1\}$

As usual we get an LP by relaxing the integrality constraints.



#### **Dual Linear Program**

max		$\sum_{j\in D} v_j$		
s.t.	$\forall i \in F$	$\sum_{j\in D} w_{ij}$	$\leq$	$f_i$
	$\forall i \in F, j \in D$	$v_j - w_{ij}$	$\leq$	$c_{ij}$
	$\forall i \in F, j \in D$	$w_{ij}$	$\geq$	0



22 Facility Location

▲ 個 ▶ ▲ 필 ▶ ▲ 필 ▶ 478/491

#### **Definition 2**

Given an LP solution  $(x^*, y^*)$  we say that facility *i* neighbours client *j* if  $x_{ij} > 0$ . Let  $N(j) = \{i \in F : x_{ij}^* > 0\}$ .



#### Lemma 3

If  $(x^*, y^*)$  is an optimal solution to the facility location LP and  $(v^*, w^*)$  is an optimal dual solution, then  $x_{ij}^* > 0$  implies  $c_{ij} \le v_j^*$ .

Follows from slackness conditions.



# Suppose we open set $S \subseteq F$ of facilities s.t. for all clients we have $S \cap N(j) \neq \emptyset$ .

Then every client j has a facility i s.t. assignment cost for this client is at most  $c_{ij} \leq v_j^*$ .

Hence, the total assignment cost is

$$\sum_j c_{i_j j} \leq \sum_j v_j^* \leq ext{OPT}$$
 ,

where  $i_j$  is the facility that client j is assigned to.



Suppose we open set  $S \subseteq F$  of facilities s.t. for all clients we have  $S \cap N(j) \neq \emptyset$ .

Then every client j has a facility i s.t. assignment cost for this client is at most  $c_{ij} \leq v_j^*$ .

nence, the total assignment cost is

 $\sum_j c_{i_j j} \leq \sum_j v_j^* \leq \mathrm{OPT}$  ,

where  $i_j$  is the facility that client j is assigned to.



Suppose we open set  $S \subseteq F$  of facilities s.t. for all clients we have  $S \cap N(j) \neq \emptyset$ .

Then every client j has a facility i s.t. assignment cost for this client is at most  $c_{ij} \le v_j^*$ .

Hence, the total assignment cost is

$$\sum_{j} c_{i_j j} \leq \sum_{j} v_j^* \leq \text{OPT}$$
 ,

where  $i_j$  is the facility that client j is assigned to.



#### Problem: Facility cost may be huge!

Suppose we can partition a subset  $F' \subseteq F$  of facilities into neighbour sets of some clients. I.e.

 $F' = \biguplus_k N(j_k)$ 

where  $j_1, j_2, \ldots$  form a subset of the clients.



### Problem: Facility cost may be huge!

Suppose we can partition a subset  $F' \subseteq F$  of facilities into neighbour sets of some clients. I.e.

 $F' = \biguplus_k N(j_k)$ 

where  $j_1, j_2, \ldots$  form a subset of the clients.



We have

 $f_{i_k}$ 



We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x^*_{ij_k}$$



We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x^*_{ij_k} \le \sum_{i \in N(j_k)} f_i x^*_{ij_k}$$



We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i y_i^* .$$



We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i y_i^* .$$

Summing over all k gives

 $\sum_{k} f_{i_k}$ 



We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i y_i^* .$$

Summing over all k gives

$$\sum_{k} f_{i_k} \le \sum_{k} \sum_{i \in N(j_k)} f_i \mathcal{Y}_i^*$$



We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i y_i^* .$$

Summing over all k gives

$$\sum_{k} f_{i_k} \leq \sum_{k} \sum_{i \in N(j_k)} f_i \mathcal{Y}_i^* = \sum_{i \in F'} f_i \mathcal{Y}_i^*$$



22 Facility Location

▲ 個 ▶ ▲ ■ ▶ ▲ ■ ▶ 483/491

We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i y_i^* .$$

Summing over all k gives

$$\sum_{k} f_{i_k} \leq \sum_{k} \sum_{i \in N(j_k)} f_i \mathcal{Y}_i^* = \sum_{i \in F'} f_i \mathcal{Y}_i^* \leq \sum_{i \in F} f_i \mathcal{Y}_i^*$$



We have

$$f_{i_k} = f_{i_k} \sum_{i \in N(j_k)} x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i x_{ij_k}^* \le \sum_{i \in N(j_k)} f_i y_i^* .$$

Summing over all k gives

$$\sum_{k} f_{i_k} \leq \sum_{k} \sum_{i \in N(j_k)} f_i \mathcal{Y}_i^* = \sum_{i \in F'} f_i \mathcal{Y}_i^* \leq \sum_{i \in F} f_i \mathcal{Y}_i^*$$

Facility cost is at most the facility cost in an optimum solution.



## Problem: so far clients $j_1, j_2, \ldots$ have a neighboring facility. What about the others?

**Definition 4** Let  $N^2(j)$  denote all neighboring clients of the neighboring facilities of client *j*.

Note that N(j) is a set of facilities while  $N^2(j)$  is a set of clients.



## Problem: so far clients $j_1, j_2, \ldots$ have a neighboring facility. What about the others?

#### **Definition 4**

Let  $N^2(j)$  denote all neighboring clients of the neighboring facilities of client j.

Note that N(j) is a set of facilities while  $N^2(j)$  is a set of clients.



Problem: so far clients  $j_1, j_2, \ldots$  have a neighboring facility. What about the others?

#### **Definition 4**

Let  $N^2(j)$  denote all neighboring clients of the neighboring facilities of client j.

Note that N(j) is a set of facilities while  $N^2(j)$  is a set of clients.









22 Facility Location

▲ 個 ▶ ▲ 필 ▶ ▲ 필 ▶ 486/491

#### Total assignment cost:

Fix k; set  $j = j_k$  and  $i = i_k$ . We know that  $c_{ij} \le v_i^*$ .



#### Total assignment cost:

- Fix k; set  $j = j_k$  and  $i = i_k$ . We know that  $c_{ij} \le v_j^*$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).



#### Total assignment cost:

- Fix k; set  $j = j_k$  and  $i = i_k$ . We know that  $c_{ij} \le v_j^*$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).





#### Total assignment cost:

- Fix k; set  $j = j_k$  and  $i = i_k$ . We know that  $c_{ij} \le v_j^*$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).

 $c_{i\ell} \leq c_{ij} + c_{hj} + c_{h\ell}$ 



#### Total assignment cost:

- Fix k; set  $j = j_k$  and  $i = i_k$ . We know that  $c_{ij} \le v_j^*$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).

$$c_{i\ell} \leq c_{ij} + c_{hj} + c_{h\ell} \leq v_j^* + v_j^* + v_\ell^*$$



#### Total assignment cost:

- Fix k; set  $j = j_k$  and  $i = i_k$ . We know that  $c_{ij} \le v_j^*$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).

$$c_{i\ell} \leq c_{ij} + c_{hj} + c_{h\ell} \leq v_j^* + v_j^* + v_\ell^* \leq 3v_\ell^*$$



#### Total assignment cost:

- Fix k; set  $j = j_k$  and  $i = i_k$ . We know that  $c_{ij} \le v_j^*$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).

$$c_{i\ell} \leq c_{ij} + c_{hj} + c_{h\ell} \leq v_j^* + v_j^* + v_\ell^* \leq 3v_\ell^*$$

Summing this over all facilities gives that the total assignment cost is at most  $3 \cdot OPT$ . Hence, we get a 4-approximation.



In the above analysis we use the inequality

$$\sum_{i\in F} f_i \mathcal{Y}_i^* \leq \text{OPT} \ .$$



22 Facility Location

◆日 → < 日 → < 日 → 487/491 In the above analysis we use the inequality

 $\sum_{i\in F} f_i \mathcal{Y}_i^* \leq \text{OPT} \ .$ 

We know something stronger namely

$$\sum_{i\in F} f_i \gamma_i^* + \sum_{i\in F} \sum_{j\in D} c_{ij} x_{ij}^* \leq \text{OPT} .$$



#### **Observation:**

Suppose when choosing a client j<sub>k</sub>, instead of opening the cheapest facility in its neighborhood we choose a random facility according to x<sup>\*</sup><sub>i,i</sub>.

Then we incur connection cost

 $\sum_{i} c_{ij_k} x_{ij_k}^*$ 

for client  $j_k.$  (In the previous algorithm we estimated this by  $\upsilon_{j_k}^*$  ).

Define

$$C_j^* = \sum_i c_{ij} x_{ij}^*$$

to be the connection cost for client j.



22 Facility Location

◆ 個 ▶ ◆ 臣 ▶ ◆ 臣 ▶ 488/491

#### **Observation:**

- Suppose when choosing a client j<sub>k</sub>, instead of opening the cheapest facility in its neighborhood we choose a random facility according to x<sup>\*</sup><sub>iik</sub>.
- Then we incur connection cost

$$\sum_{i} c_{ij_k} x^*_{ij_k}$$

for client  $j_k.$  (In the previous algorithm we estimated this by  $\boldsymbol{v}_{j_k}^*$  ).

Define

$$C_j^* = \sum_i c_{ij} x_{ij}^*$$

to be the connection cost for client j.



#### **Observation:**

- Suppose when choosing a client j<sub>k</sub>, instead of opening the cheapest facility in its neighborhood we choose a random facility according to x<sup>\*</sup><sub>iik</sub>.
- Then we incur connection cost

$$\sum_{i} c_{ij_k} x^*_{ij_k}$$

for client  $j_k$ . (In the previous algorithm we estimated this by  $v_{j_k}^*$ ).

Define

$$C_j^* = \sum_i c_{ij} x_{ij}^*$$

to be the connection cost for client j.


We only try to open a facility once (when it is in neighborhood of some  $j_k$ ). (recall that neighborhoods of different  $j'_k s$  are disjoint).

We open facility i with probability  $x_{ij_k} \le y_i$  (in case it is in some neighborhood; otw. we open it with probability zero).

Hence, the expected facility cost is at most

 $\sum_{i\in F}f_i\mathcal{Y}_i \cdot$ 



We only try to open a facility once (when it is in neighborhood of some  $j_k$ ). (recall that neighborhoods of different  $j'_k s$  are disjoint).

We open facility i with probability  $x_{ij_k} \le y_i$  (in case it is in some neighborhood; otw. we open it with probability zero).

Hence, the expected facility cost is at most

 $\sum_{i\in F} f_i \mathcal{Y}_i \cdot$ 



We only try to open a facility once (when it is in neighborhood of some  $j_k$ ). (recall that neighborhoods of different  $j'_k s$  are disjoint).

We open facility i with probability  $x_{ij_k} \leq y_i$  (in case it is in some neighborhood; otw. we open it with probability zero).

Hence, the expected facility cost is at most

 $\sum_{i\in F} f_i \gamma_i$  .



We only try to open a facility once (when it is in neighborhood of some  $j_k$ ). (recall that neighborhoods of different  $j'_k s$  are disjoint).

We open facility *i* with probability  $x_{ij_k} \le y_i$  (in case it is in some neighborhood; otw. we open it with probability zero).

Hence, the expected facility cost is at most

 $\sum_{i\in F}f_i\mathcal{Y}_i$  .







- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- ► If we assign a client l to the same facility as i we pay at most

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2v_j^* \le \sum_{j} C_j^* + 2\text{OPT}$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2v_j^* \le \sum_{j} C_j^* + 20PT$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client  $\ell$  to the same facility as i we pay at most

$$\sum_{i} c_{ij} x_{ijk}^* + c_{hj} + c_{h\ell} \le C_j^* + v_j^* + v_\ell^* \le C_\ell^* + 2v_\ell^*$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2v_j^* \le \sum_{j} C_j^* + 2\text{OPT}$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

$$\sum_{i} c_{ij} \chi_{ij_{k}}^{*} + c_{hj} + c_{h\ell} \le C_{j}^{*} + v_{j}^{*} + v_{\ell}^{*} \le C_{\ell}^{*} + 2v_{\ell}^{*}$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2\nu_j^* \le \sum_{j} C_j^* + 20PT$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

$$\sum_{i} c_{ij} \chi_{ij_k}^* + c_{hj} + c_{h\ell} \le C_j^* + \nu_j^* + \nu_\ell^* \le C_\ell^* + 2\nu_\ell^*$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2\nu_j^* \le \sum_{j} C_j^* + 20PT$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

$$\sum_{i} c_{ij} x_{ij_k}^* + c_{hj} + c_{h\ell} \le C_j^* + v_j^* + v_\ell^* \le C_\ell^* + 2v_\ell^*$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_{j}^{*} + \sum_{j} 2v_{j}^{*} \le \sum_{j} C_{j}^{*} + 2\text{OPT}$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

$$\sum_{i} c_{ij} x_{ij_k}^* + c_{hj} + c_{h\ell} \le C_j^* + v_j^* + v_\ell^* \le C_\ell^* + 2v_\ell^*$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2v_j^* \le \sum_{j} C_j^* + 2\text{OPT}$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

$$\sum_{i} c_{ij} x_{ij_k}^* + c_{hj} + c_{h\ell} \le C_j^* + v_j^* + v_\ell^* \le C_\ell^* + 2v_\ell^*$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2\nu_j^* \le \sum_{j} C_j^* + 2\text{OPT}$$

Hence, it is at most 20PT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

$$\sum_{i} c_{ij} x_{ij_k}^* + c_{hj} + c_{h\ell} \le C_j^* + v_j^* + v_\ell^* \le C_\ell^* + 2v_\ell^*$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_j^* + \sum_{j} 2\nu_j^* \le \sum_{j} C_j^* + 2\text{OPT}$$

Hence, it is at most 2OPT plus the total assignment cost in an optimum solution.

- Fix k; set  $j = j_k$ .
- Let  $\ell \in N^2(j)$  and h (one of) its neighbour(s) in N(j).
- If we assign a client l to the same facility as i we pay at most

$$\sum_{i} c_{ij} x_{ij_k}^* + c_{hj} + c_{h\ell} \le C_j^* + v_j^* + v_\ell^* \le C_\ell^* + 2v_\ell^*$$

Summing this over all clients gives that the total assignment cost is at most

$$\sum_{j} C_{j}^{*} + \sum_{j} 2\nu_{j}^{*} \le \sum_{j} C_{j}^{*} + 2\text{OPT}$$

Hence, it is at most 2OPT plus the total assignment cost in an optimum solution.