A crucial ingredient for the design and analysis of approximation algorithms is a technique to obtain an upper bound (for maximization problems) or a lower bound (for minimization problems).

Therefore Linear Programs or Integer Linear Programs play a vital role in the design of many approximation algorithms.

Many important combinatorial optimization problems can be formulated in the form of an Integer Program.

Note that solving Integer Programs in general is **NP-complete!**

EADS II ||||||| © Harald Räcke

272

270

Definition 2

An Integer Linear Program or Integer Program is a Linear Program in which all variables are required to be integral.

Definition 3

A Mixed Integer Program is a Linear Program in which a subset of the variables are required to be integral.

EADS II © Harald Räcke

12 Integer Programs



271





IP-Formulation of Set Cover						
	min		$\sum_i w_i x_i$)
		$\forall u \in U$	$\sum_{i:u\in S_i} x_i$	\geq	1	
		$\forall i \in \{1, \ldots, k\}$	x_i	\geq	0	
		$\forall i \in \{1, \dots, k\}$	x_i	integral		
50 00 EADS II		12 Integ	er Programs			
	äcke	12 mcg				275



Maximum Weighted Matching

Given a graph G = (V, E), and a weight w_e for every edge $e \in E$. Find a subset of edges of maximum weight such that no vertex is incident to more than one edge.

max		$\sum_{e\in E} w_e x_e$		
s.t.	$\forall v \in V$	$\sum_{e:v \in e} x_e$	\leq	1
	$\forall e \in E$	x_e	\in	$\{0, 1\}$

EADS II © Harald Räcke	12 Integer Programs	278

Knapsack

Given a set of items $\{1, \ldots, n\}$, where the *i*-th item has weight w_i and profit p_i , and given a threshold K. Find a subset $I \subseteq \{1, \dots, n\}$ of items of total weight at most K such that the profit is maximized.

max		$\sum_{i=1}^{n} p_i x_i$		
s.t.		$\sum_{i=1}^{n} w_i x_i$	\leq	Κ
	$\forall i \in \{1, \dots, n\}$	x_i	\in	$\{0, 1\}$

Maximum Independent Set

Given a graph G = (V, E), and a weight w_v for every node $v \in V$. Find a subset $S \subseteq V$ of nodes of maximum weight such that no two vertices in *S* are adjacent.

max		$\sum_{v \in V} w_v x_v$		
s.t.	$\forall e = (i, j) \in E$	$x_i + x_j$	\leq	1
	$\forall v \in V$	x_v	\in	$\{0, 1\}$

EADS II © Harald Räcke	12 Integer Programs	279



A linear program LP is a relaxation of an integer program IP if any feasible solution for IP is also feasible for LP and if the objective values of these solutions are identical in both programs.

We obtain a relaxation for all examples by writing $x_i \in [0, 1]$ instead of $x_i \in \{0, 1\}$.

EADS II © Harald Räcke

By solving a relaxation we obtain an upper bound for a maximization problem and a lower bound for a minimization problem.

	12 Integer Programs	
🛛 💾 🛛 🖓 © Harald Räcke		282



